

This leads to a wrong updating result. Accordingly, it affirms the weakness of worldview-2 elevation model which produced by image matching of worldview-2 satellite stereo images comparing LiDAR DSM.

ACKNOWLEDGEMENT

The authors are grateful from DLR for giving data.

REFERENCES

- E. Hanson and E. Wolff, "Change detection for update of topographic databases through multi-level region-based classification of VHR optical and SAR data," *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, vol. 38, 2010.
- E. Hanson and E. Wolff, "Change detection for update of topographic databases through multi-level region-based classification of VHR optical and SAR data," *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, vol. 38, 2010.
- T. T. Vu, M. Matsuoka, and F. Yamazaki, "LIDAR-based change detection of buildings in dense urban areas," in *Geoscience and Remote Sensing Symposium, 2004. IGARSS'04. Proceedings. 2004 IEEE International*, 2004, vol. 5, pp. 3413–3416.
- Krauss, T., Sirmacek, B., Arefi, H., & Reinartz, P. (2012). Fusing stereo and multispectral data from WorldView-2 for urban modeling. In *Proc. of SPIE Vol (Vol. 8390, p. 83901X–1)*
- J. Tian, H. Chaabouni-Chouayakh, and P. Reinartz, "3D building change detection from high resolution spaceborne stereo imagery," in *Multi-Platform/Multi-Sensor Remote Sensing and Mapping (M2RSM), 2011 International Workshop on*, 2011, pp. 1–7.
- Gruen, A., & Akca, D. (2005). Least squares 3D surface and curve matching. *ISPRS Journal of Photogrammetry and Remote Sensing*, 59(3), 151–174.
- Arefi, H., & Hahn, M. (2005). A hierarchical procedure for segmentation and classification of airborne LIDAR images. In *INTERNATIONAL GEOSCIENCE AND REMOTE SENSING SYMPOSIUM (Vol. 7, p. 4950)*
- K. Choi, I. Lee, and S. Kim, "A feature based approach to automatic change detection from LiDAR data in Urban areas," *Laserscanning09*, pp. 259–264, 2009
- J. Trinder and M. Salah, "Aerial images and LiDAR data fusion for disaster change detection," in *The XXII Congress of the International Society for Photogrammetry and Remote Sensing*, Melbourne, 2012, pp. 227–232.
- J. Tian, H. Chaabouni-Chouayakh, P. Reinartz, T. Krauss, and P. d' Angelo, "Automatic 3D change detection based on optical satellite stereo imagery," in *ISPRS TC VII Symposium*, Vienna, 2010.
- L. Zhu, H. Shimamura, K. Tachibana, Y. Li, and P. Gong, "Building change detection based on object extraction in dense urban areas," *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci. Beijing China*, vol. 37, pp. 905–908.
- J. Tian, P. Reinartz, and P. d' Angelo, "Change Detection Analysis of Forest Areas Using Satellite Stereo Data.," in *GIL Jahrestagung*, 2012, pp. 311–314.
- G. Vosselman, B. G. H. Gorte, and G. Sithole, "Change detection for updating medium scale maps using laser altimetry," *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, vol. 34, pp. 1–6, 2004.
- J. Zhang, Z. Zhang, H. FAN, Z. FANG, and Z. LIU, "Change Detection in Urban Area Based on Stereo Image Pairs from Different Duration," *Int. Arch. Photogramm. REMOTE Sens.*, vol. 33, no. B4/3; PART 4, pp. 1203–1208, 2000.
- M. Rutzinger, B. Rűf, B. Höfle, and M. Vetter, "Change detection of building footprints from airborne laser scanning acquired in short time intervals," in *ISPRS TC VII Symposium*, 2010, vol. 100, pp. 475–480.
- H. Murakami, K. Nakagawa, H. Hasegawa, T. Shibata, and E. Iwanami, "Change detection of buildings using an airborne laser scanner," *ISPRS J. Photogramm. Remote Sens.*, vol. 54, no. 2, pp. 148–152, 1999.
- M. Rehor, "Classification of building damage based on laser scanning data," *Photogramm. J. Finl.*, vol. 20, no. 2, 2007.
- F. Jung, "Detecting building changes from multitemporal aerial stereopairs," *ISPRS J. Photogramm. Remote Sens.*, vol. 58, no. 3, pp. 187–201, 2004.
- T. Vögtle and E. Steinle, "Detection and recognition of changes in building geometry derived from multitemporal laserscanning data," *Int. Arch. Photogramm. Remote Sens.*, vol. 35, no. B2, pp. 428–433, 2004.
- H. Murakami, K. Nakagawa, T. Shibata, and E. Iwanami, "Potential of an airborne laser scanner system for change detection of urban features and orthoimage development," *Int. Arch. Photogramm. Remote Sens.*, vol. 32, pp. 422–427, 1998.
- T. Santos, S. Freire, A. Fonseca, and J. A. Tenedório, "Producing a building change map for urban management," *EARSeL EProceedings*, vol. 10, no. 1, p. 56, 2011.
- J. Tian, J. Leitloff, T. Krauss, and P. Reinartz, "Region based forest change detection from Cartosat-1 stereo imagery," in *ISPRS Hannover Workshop*, 2011.
- H. Chaabouni-Chouayakh, T. Krauss, P. d'Angelo and P. Reinartz, "3D change detection inside urban areas using different Digital Surface Models," In: *PapadotisN., Pierrot-Deseil li gny M., Mal let C., Tournai reO. (Eds), IAPRS, Vol. XXXVII I, Part 3B – Saint-Mandé, France, September 1-3, 2010*
- P. Wezyk, M. Szostak, P. Tompalski, and G. Zajaczkowski, "The role of Airborne Laser scanning in updating and revising GIS databases-a case study in the Polish State Forest (Chojna District)," *Proc. Silvilaser*, 2010.
- M. Rezaee, "Object Based Classification of High Spatial Resolution WorldView-II Imagery in Urban Area. " *M.Sc. Thesis*, university of Tehran. Iran, 2012.