















- Environment, 100, pp. 474-485.
- [10] Chen, J. M.; Pavlic, G.; Brown, L.; Cihlar, J.; Leblanc, S.G.; White, H.P.; Hall, R.J.; Peddle, D.; King, D.J.; Trofymow, J.A.; Swift, E.; Vander, S.J.; Pellikka, P., 2002. Validation of Canada-wide leaf area index maps using ground measurements and high and moderate resolution satellite imagery. *Remote Sensing of Environment*, 80, pp. 165-184.
- [11] Sandmeier, S.R.; Deering, D.W., A new approach to derive canopy structure information for boreal forests using spectral BRDF data. *IEEE IGARSS 1999 Proceedings Germany: Hamburg*.
- [12] Sandmeier, S.R.; Miller, C.; Hosgood, B.; Andreoli, G., 1998. Physical mechanisms in hyperspectral BRDF data of grass and watercress. *Remote Sensing of Environment*, 66, pp. 222-233.
- [13] Lacaze, R.; Chen, J.M.; Roujean, J.L.; Leblanc, S.G., 2002. Retrieval of vegetation clumping index using hot spot signatures measured by POLDER instrument. *Remote Sensing of Environment*, 79, pp. 84-95.
- [14] Pocewicz, A.; Vierling, L.A.; Lentile, L.B.; Smith, R., 2007. View angle effects on relationships between MISR vegetation indices and leaf area index in a recently burned ponderosa pine forest. *Remote Sensing of Environment*, 107, pp. 322-333.
- [15] Wu, C.; Niu, Z.; Wang, J.; Gao, S.; Huang, W., 2010. Predicting leaf area index in wheat using angular vegetation indices derived from in situ canopy measurements. *Canadian Journal of Remote Sensing*, 36(4), pp. 301-312.
- [16] Kuusk, A., 2001. A two-layer canopy reflectance model. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 71, pp. 1-9.
- [17] Liu, L.Y.; Wang, J.H.; Bao, Y.S.; Huang, W.J.; Ma, Z.; Zhao, H., 2006. Predicting winter wheat condition, grain yield and protein content using multi-temporal EnviSat-ASAR and Landsat TM satellite images. *International Journal of Remote Sensing*, 27, pp. 737-753.
- [18] Verrelst, J.; Schaepman, M.E.; Koetz, B.; Kneubuehler, M., 2007. Angular sensitivity analysis of vegetation indices derived from CHRIS/PROBA data, *Remote Sensing of Environment*, 112(5), pp. 2341-2353.
- [19] Fan, W.J.; Xu, X.R.; Liu, X.C.; Yan, B.Y.; Cui, Y.K., 2010. Accurate LAI retrieval method based on PROBA/CHRIS data. *Hydrology and Earth System Sciences*, 14, pp. 1499-1507.
- [20] CHRIS/PROBA Website: <http://earth.esa.int/missions/thirdpartymission/proba.html>.
- [21] Gao, S.; Niu, Z.; Wu, C., 2010. Multi-polarization Envisat-ASAR images as a function of leaf area index (LAI) of White Poplar and Desert Date plantations. *International Journal of Remote Sensing*, 31(4), pp. 1095-1102.
- [22] Li, X.; Li, X.W.; Li, Z.Y.; Ma, M.G.; Wang, J.; Xiao, Q.; Liu, Q.; Che, T.; Chen, E.X.; Yan, G.J.; Hu, Z.Y.; Zhang, L.X.; Chu, R.Z.; Su, P.X.; Liu, Q.H.; Liu, S.M.; Wang, J.D.; Niu, Z.; Chen, Y.; Jin, R.; Wang, W.Z.; Ran, Y.H.; Xin, X.Z.; Ren, H.Z., 2009. Watershed Allied Telemetry Experimental Research, *Journal of Geophysical Research*, 114(D22103).
- [23] Fang, H.; Liang, S.; Kuusk, A., 2003. Retrieving leaf area index using a genetic algorithm with a canopy radiative transfer model. *Remote Sensing of Environment* 85(3), pp. 257-270.
- [24] Hosgood, B.; Jacquemoud, S.; Andreoli, G.; Verdebout, J.; Pedrini, J.; Schmuck, G., 1995. Leaf Optical Properties Experiment 93 (LOPEX 93), Report EUR-16095-EN, European Commission, Joint Research Center, Institute for Remote Sensing Applications, Ispra, Italy.
- [25] Houborg, R.; Soegaard, H.; Boegh, E., 2007. Combining vegetation index and model inversion methods for the extraction of key vegetation biophysical parameters using Terra and Aqua MODIS reflectance data. *Remote Sensing of Environment*, 106, pp. 39-58.
- [26] Houborg, R.; Boegh E., 2008. Mapping leaf chlorophyll and leaf area index using inverse and forward canopy reflectance modeling and SPOT reflectance data. *Remote Sensing of Environment*, 112(1), pp. 186-202.
- [27] Xiao, Z.; Liang, S.; Wang, J.; Jiang, B.; Li, X., 2011. Real-time retrieval of leaf area index from MODIS time series data. *Remote Sensing of Environment*, 115(1), pp. 97-106.
- [28] Kuusk, A., 2009. Two-layer Canopy Reflectance Model ACRM User Guide.
- [29] Houborg, R.; Anderson, M.; Daughtry, C., 2009. Utility of an image-based canopy reflectance modeling tool for remote estimation of LAI and leaf chlorophyll content at the field scale. *Remote Sensing of Environment*, 113(1), pp. 259-274.
- [30] Price, J.C., On the information content of soil reflectance spectral, 1990. *Remote Sensing of Environment*, 33(2), pp. 113-121.
- [31] Rouse, J.W.J.; Haas, R.H.; Schell, J.A.; Deering, D.W., 1974. Monitoring vegetation systems in the Great Plains with ERTS. In: *Third ERTS Symposium*, NASA SP-351, Washington DC, pp. 309-317.
- [32] Li, X.; Gao, F.; Wang, J.; Yang, H., 2002. Bi-directional normalized difference vegetation index. Concept and application, *Progress in Natural Science*, 12 (2), pp. 115-119.
- [33] Wu, C.; Niu, Z.; Tang, Q.; Huang, W., Estimating chlorophyll content from hyperspectral vegetation indices: Modeling and validation. *Agricultural and Forest Meteorology* 2008, 148, pp. 1230-1241.
- [34] Haboudane, D.; John, R.; Millera, J.R.; Tremblay, N.; Zarco-tejada, P.J.; Dextraze, L., 2002. Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture. *Remote Sensing of Environment*, 81, pp. 416-426.
- [35] Fang, H.; Wei, S.; Liang, S., 2012. Validation of MODIS and CYCLOPES LAI products using global field measurement data. *Remote Sensing of Environment*, 119, pp. 43-54.
- [36] Wang Lijuan, Niu Zheng, Hou Xuehui, Gao Shuai, 2013. The study of LAI estimation using a new vegetation index based on CHRIS data. *Spectroscopy and Spectral Analysis*, 33(4), pp: 1082-1086.