

cloud data is large, so the efficient data processing method should be studied. Point cloud contains a great deal of information, the way to dig out more information and make full use of it needs to be considered.

REFERENCES

- Alonzo, M., Bookhagen, B., Roberts, D. A., 2014. Urban tree species mapping using hyperspectral and lidar data fusion. *Remote Sensing of Environment*, 148, 70-83.
- Bright, B. C., Hicke, J. A., Hudak, A. T., 2012. Estimating aboveground carbon stocks of a forest affected by mountain pine beetle in Idaho using lidar and multispectral imagery. *Remote Sensing of Environment*, 124, 270-281.
- Estornell, J., Ruiz, L. A., Velázquez-Martí B., López-Cortés, I., Salazar, D., Fernández-Sarrá, A., 2015. Estimation of pruning biomass of olive trees using airborne discrete-return LiDAR data. *Biomass and Bioenergy*, 81, 315-321.
- Falkowski, M. J., Smith, A. M., Hudak, A. T., Gessler, P. E., Vierling, L. A., Crookston, N. L., 2006. Automated estimation of individual conifer tree height and crown diameter via two-dimensional spatial wavelet analysis of lidar data. *Canadian Journal of Remote Sensing*, 32(2), 153-161.
- Fang, L., Yang, B., 2013. Automated extracting structural roads from mobile laser scanning point clouds. *Acta Geodaetica et Cartographica Sinica*, 42(2), 260-267.
- Fitzgibbon, A., Pilu, M., Fisher, R. B., 1999. Direct least square fitting of ellipses. *IEEE Transactions on pattern analysis and machine intelligence*, 21(5), 476-480.
- Huang, C., Wen, S., Ouyang, Q., Yang, L., 2015. Research on biomass and carbon storage of main tree species in Wangcheng district of Changsha city. *Journal of Central South University of Forestry & Technology*, 35(9), 128-132.
- Maas, H. G., Bienert, A., Scheller, S., Keane, E., 2008. Automatic forest inventory parameter determination from terrestrial laser scanner data. *International journal of remote sensing*, 29(5), 1579-1593.
- Ministry of Agriculture and Forestry of P.R. China, 1978. *Standing Volume Tables*. Standards Press of China, Beijing, 97.
- Shi, X., 2010. Preliminary study on fixing carbon dioxide releasing oxygen and retention of dust of street trees in Beijing. Beijing: Beijing Forestry University, 36 (Master's thesis).
- Sirmacek, B., Lindenbergh, R., 2015. Automatic classification of trees from laser scanning point clouds. ISPRS.
- Wang, B., Liu, M., Zhang, B., 2009. Dynamics of net production of Chinese forest vegetation based on forest inventory data. *Forest Resources Management*, 1(3), 35-43.
- Wang, Y., Zhou, G., Jiang, Y., Yang, Z., 2001. Estimating biomass and NPP of larch forests using forest inventory data. *Chinese Journal of Plant Ecology*, 25(4), 420-425.
- Wei, Z., Yang, B., Li, Q., 2012. Automatic extraction of building footprints from mobile LIDAR point clouds. *Journal of Remote Sensing*, 16(2), 286-296.
- Wu, B., Yu, B., Yue, W., Shu, S., Tan, W., Hu, C., Huang, Y., Wu, J., Liu, H., 2013. A voxel-based method for automated identification and morphological parameters estimation of individual street trees from mobile laser scanning data. *Remote Sensing*, 5(2), 584-611.
- Yang, B., Wei, Z., Li, Q., Li, J., 2012. Automated extraction of street-scene objects from mobile lidar point clouds. *International Journal of Remote Sensing*, 33(18), 5839-5861.
- Yang, S., Li, Y., Li, K., Mao, J., 2014. Tree extraction from vehicle-borne LiDAR data. *Engineering of Surveying and Mapping*, 23(8), 23-47.
- Yu, X., Liang, X., Hyypä J., Kankare, V., Vastaranta, M., Holopainen, M., 2013. Stem biomass estimation based on stem reconstruction from terrestrial laser scanning point clouds. *Remote sensing letters*, 4(4), 344-353.